

Effect of zinc and magnesium oxide phytonanoparticles on the production of rebaudiosides and phenolic compounds in *Stevia rebaudiana* Bertoni plants

Efecto de las fitonanopartículas de óxido de zinc y magnesio en la producción de rebaudiósidos y compuestos fenólicos en plantas de *Stevia rebaudiana* Bertoni

María Concepción Velázquez-Gamboa¹✉, Ludwi Rodríguez-Hernández²✉, Miguel Abud-Archila¹✉, Federico Antonio Gutiérrez-Miceli¹✉, Víctor Manuel Ruíz-Valdiviezo¹✉, Benjamín Valdez-Salas³✉, Dalia Margarita Ferrer-Sánchez¹✉, María Celina Luján-Hidalgo^{2*}✉

¹ Tecnológico Nacional de México/IT de Tuxtla Gutiérrez, División de Estudios de Posgrado e Investigación, Carretera Panamericana km. 1080, Tuxtla Gutiérrez, Chiapas, C.P. 29050, México.

² Instituto Tecnológico Superior de Cintalapa, Carretera Panamericana km. 995 C.P. 30400 Cintalapa, Chiapas, México.

³ Universidad Autónoma de Baja California, Instituto de Ingeniería, Mexicali, Baja California, C. P. 21100, México.

ABSTRACT

The demand for natural sweeteners has been increasing. This trend is influenced by several factors, including concerns and controversies surrounding synthetic sweeteners. This study aimed to determine the effect of elicitation with ZnO and MgO phytonanoparticles (PTNPs) on *Stevia rebaudiana* plants. The synthesis of PTNPs is possible due to reduction potential of the natural aqueous extract derived from Moringa oleifera leaves. The characterization of PTNPs was carried out using spectroscopic and microscopic methods. Scanning electron microscope images of the ZnO PTNPs showed the presence of agglomerated rod-shaped nanoparticles, while the MgO PTNPs were irregularly shaped and agglomerated. Dynamic light scattering analysis showed an average size of 9 - 30 nm for ZnO PTNPs and 5 - 10 nm for MgO PTNPs, while zeta potential (ζ) values validated the stability of the PTNPs. The infrared spectra revealed bands corresponding to stretching vibrations of the bonds between the groups involved in the reduction and stabilization of PTNPs. The highest production of rebaudiosides A, E and M was obtained using 1 mg/L of ZnO PTNPs. The elicitation with PTNPs, mainly with ZnO at the concentration of 20 mg/L, induces the production of molecules with antioxidant capacity, obtaining a 137 % increase of chlorogenic acid, while that of caffeic acid is 185 %. These results indicated that the application of ZnO and MgO PTNPs induced a higher content of antioxidant compounds in *Stevia rebaudiana*.

Keywords: Elicitation, Rebaudioside M, Antioxidant compounds, Chlorogenic acid.

RESUMEN

La demanda de edulcorantes naturales ha ido en aumento; Esta tendencia está influenciada por varios factores, incluidas las preocupaciones y controversias en torno a los edulcorantes sintéticos. Este estudio tiene como objetivo determinar el efecto de la estimulación con fitonanopartículas de ZnO y MgO (PTNPs) de plantas de *Stevia rebaudiana*. La síntesis de PTNPs es posible gracias al potencial de reducción del

extracto acuoso natural derivado de las hojas de Moringa oleifera. La caracterización de las PTNPs se realizó mediante métodos espectroscópicos y microscópicos. Las imágenes de microscopio electrónico de barrido de las PTNPs de ZnO mostraron la presencia de nanopartículas aglomeradas en forma de varilla, mientras que las PTNPs de MgO tenían forma irregular y estaban aglomeradas. El análisis de dispersión dinámica de la luz mostró un tamaño promedio de 9 a 30 nm para los PTNPs de ZnO y de 5 a 10 nm para los PTNPs de MgO, mientras que los valores del potencial zeta (ζ) validaron la estabilidad de los PTNPs. Los espectros infrarrojos revelaron bandas correspondientes a vibraciones de estiramiento de los enlaces de los grupos implicados en la reducción y estabilización de los PTNPs. La mayor producción de rebaudiósidos A, E y M se obtuvo utilizando 1 mg/L de ZnO PTNP. La elicitación con PTNPs principalmente con ZnO a partir de la concentración de 20 mg/L induce la producción de moléculas con capacidad antioxidante, obteniendo un aumento del 137 % del ácido clorogénico, mientras que el del ácido cafeico es del 185 %. Estos resultados indicaron que la aplicación de PTNPs de ZnO y MgO indujo un mayor contenido de compuestos antioxidantes en *Stevia rebaudiana*.

Palabras clave: Estimulación, Rebaudiósido M, Compuestos antioxidantes, Ácido clorogénico

INTRODUCTION

At the commercial level, there has been a recent increase in consumer interest for the use of substances of natural origin, especially plant extracts with therapeutic properties, to treat diseases such as obesity, kidney disease, cardiovascular disease, cancer, etc., among which is the stevia plant (Saraiva et al., 2020).

Stevia rebaudiana Bertoni is a plant of commercial interest, known for its high content of steviol glycosides (SGs) (Cauich-Cauich et al., 2022; Javed et al., 2017a). In 2024, the global market for *S. rebaudiana* will be valued at \$405.6 million and is projected to reach \$739.4 million in 2034,

*Author for correspondence: María Celina Luján Hidalgo
e-mail: maria.lh@tuxtla.tecnm.mx

Received: April 16, 2024

Accepted: August 13, 2024

Published: Septiembre 4, 2024

reflecting a compound annual growth rate (CAGR) of 6.2 % (Stevia Market, 2024).

Stevioside (Stev), rebaudioside A (Reb A) and rebaudioside C (Reb C), are the most abundant SGs in the leaves, and commercially these molecules have been attributed the sweetening power. Other SGs, such as rebaudiosides D, E, and M, have greater sweetening capacity, however, they are found at very low concentrations (Gupta *et al.*, 2013). In addition, other studies have shown that the leaves of *S. rebaudiana* contain antioxidant compounds such as ascorbic acid, gallic acid and other phenolic compounds, including flavonoids, alkaloids, chlorophylls, xanthophylls and hydroxycinnamic acids (Pandey *et al.*, 2018) that confer additional functional properties to the sweetening power, so the implementation of agronomic techniques that induce the biosynthesis of these compounds is an area of interest.

Today, the pharmaceutical and food industries have a growing interest in improving the quantity and quality of the active principles of medicinal plants (Ramezani *et al.*, 2020). Increasing the production of secondary metabolites by modifying plant metabolism, is an efficient approach that could be achieved through several means.

Elicitors are biotic or abiotic substances or factors that can induce a metabolic response in plants depending mainly on their concentration, chemical nature, size and growth stage of the plant (Wang *et al.*, 2017). Some phytohormones, such as methyl jasmonate (MeJa), have been used as elicitors. This phytohormone plays a crucial role in signaling pathways leading to the accumulation of secondary metabolites, which can have diverse applications including pharmaceuticals, flavors, and fragrances. However, their cost can be a challenge, especially for large-scale applications (Lucho *et al.*, 2018; Moharramnejad *et al.*, 2019).

Recently, research in the field of nanotechnology for signaling the effects of environmental stress on plant physiology is advancing at a rapid pace (Abdel-Aziz *et al.*, 2020). Nanotechnology has the potential to revolutionize agriculture and significantly impact food and crop production (Magdaleno-García *et al.*, 2023). Scientists are increasingly interested in employing biotechnological techniques to enhance the production of secondary metabolites and explore various applications in agriculture such as the formulation of nanopesticides and nanofertilizers (Nongbet *et al.*, 2022).

Recent studies have shown that nanoparticles can act as elicitors (Vijayakumar *et al.*, 2022) as these particles can rapidly penetrate plant cells, elevate protein levels and stimulate gene expression (Masarovicová *et al.*, 2013). Javed *et al.* (2017a) investigated the effect of different concentrations of zinc oxide chemical nanoparticles (ZnO NPs) on growth and steviol glycoside production in micropropagated shoots of *S. rebaudiana*. They found that 1 mg/L of ZnO nanoparticles produced the highest percentage of shoot formation (89.6 %), by doubling the content of stevioside and rebaudioside A compared to the control, suggesting a positive influence of nanoparticles on plant growth and glucoside production.

The green synthesis of nanoparticles is particularly

attractive due to their biocompatibility, low toxicity, and eco-friendly nature (Bouttier-Figueroa *et al.*, 2023). Key considerations for synthesizing and producing plant-based nanoparticles include the selection of environmentally friendly or non-harmful solvents for stabilization (Raliya *et al.*, 2014). Traditional chemical methods used in various industries can be costly and environmentally hazardous, especially when they involve toxic and hazardous chemicals (Krishnani *et al.*, 2022). This has driven increased interest in green chemistry and sustainable practices that aim to minimize the environmental and health impacts of chemical processes (Gowri *et al.*, 2019).

The main objective of this research was to determine how the application of ZnO and MgO nanoparticles as elicitors affects the content of certain compounds in dried *Stevia rebaudiana* leaves. Specifically, the study aims to measure changes in the levels of rebaudiosides A, D, M and address the metabolic profile of phenolic compounds such as gallic acid, chlorogenic acid, caffeic acid and caffeine.

MATERIAL AND METHODS

Synthesis and characterization of ZnO and MgO phytonanoparticles from *Moringa oleifera*

The plant extract used for the synthesis of PTNPs was obtained from previously pulverized dehydrated leaves of *Moringa oleifera*, as described by Rodríguez-Pérez *et al.* (2015). Briefly, 500 mg of plant material were mixed with 25 mL of distilled water and maintained at 25 °C for 1 hour under continuous magnetic stirring. The mixture was sonicated at 360 W (COLE-PARMER, Vernon Hills, IL, USA) for 15 minutes at 25 °C and centrifuged at 4500 rpm for 10 minutes (HERMLE, Germany) to remove solids. The extract was filtered through a 0.45 µm Millipore membrane.

The synthesis of zinc and magnesium PTNPs was carried out following the methodology described by Chaudhuri and Malodia (2017), with modifications as reported by Velázquez-Gamboa *et al.* (2021).

Characterization techniques were used to obtain a comprehensive understanding of the chemical composition, size distribution, and physical morphology of ZnO and MgO PTNPs, which were characterized by FT-IR (NICOLET, Wisconsin, USA), SEM (JEOL 6010LA), dynamic light scattering (DLS, Montgomeryville, PA, USA), and ζ potential (Microtrac, Montgomeryville PA, USA) (Selvakesavan *et al.*, 2023).

Plant cultivation (*Stevia rebaudiana* Bertoni) and elicitation with ZnO and MgO phytonanoparticles

Stevia rebaudiana (Var. Morita II) plants were grown in plastic seedbeds using a mixture of peat and agrolite (4:1 v/v). After the development of 10 true leaves, plants were transplanted into polyethylene bags (20 x 30 cm) containing the same inert substrate. The experiment was carried out in an anti-aphid house in Cintalapa de Figueroa, Chiapas, Mexico (16°45'N 93°51'W). The temperature in the anti-aphid house ranged between 29 - 32 °C and relative humidity below 70 %. Plant



nutrition was provided with modified Hoagland nutrient solution without zinc and magnesium.

At 30 days after transplantation (15 and 20 true leaves), different experimental treatments with ZnO and MgO PTNPs were applied. Plants were sprayed individually from the roots twice a week (1 mL per plant) for four weeks. The treatments evaluated were: T1 (control without PTNPs), T2 (MeJa 22 mg/L), T3 (PTNPs ZnO 1 mg/L), T4 (20 mg/L ZnO PTNPs), T5 (30 mg/L ZnO PTNPs), T6 (1 mg/L MgO PTNPs), T7 (20 mg/L MgO PTNPs), and T8 (30 mg/L MgO PTNPs). At the end of the experiment, plants were harvested and dried at 60 °C for 48 h. They were ground and stored in the dark at 4 °C until use.

***Stevia rebaudiana* extract preparation**

The method described by Gasmalla *et al.* (2017) was used for the extraction of the steviol glycosides.

Rebaudiosides A, D, E and M quantification

The quantification of SGs (rebaudioside A, D and M) present in the extract of *S. rebaudiana* Bertoni was carried out by HPLC UV-Visible (High-performance liquid chromatography with UV-Vis detector, Flexar PerkinElmer, USA), according to the methodology reported by Velázquez-Gamboa *et al.* (2021).

Gallic acid, chlorogenic acid, caffeic acid, and caffeine contents

The system used for the analysis of the various phenolic compounds consisted of a Perkin Elmer high-performance liquid chromatograph. A reversed-phase C18 column (150 x 4.4 mm) was used at a wavelength of 270 nm for gallic acid and caffeine, and 325 nm for chlorogenic acid and caffeic acid.

The mobile phases used were 0.1 % formic acid (solvent A) and methanol (solvent B). Twenty μ L of sample were injected, a solvent gradient elution program was used starting at a 15 % (A)/85 % (B), increasing to 80 % (A)/20 % (B) throughout 50 min, then decreasing to 15 % (A)/85 % (B) over 2 min, and finally maintaining the same ratios for 5 min. External standards of gallic acid, chlorogenic acid, caffeic acid, and caffeine HPLC grade (Sigma Aldrich, USA) were used, and the phenolic compounds were expressed in mg/g of dry leaf.

Statistical analysis

A completely randomized design was used, considering 21 replicas per treatment. An analysis of variance and Fisher's LSD test were performed ($p < 0.005$). Statistical analysis was performed using Statgraphics Centurion XVII software.

RESULTS AND DISCUSSION

ZnO and MgO phytonanoparticles characterization

The formation of PTNPs was evidenced by the color changes in the precursor solutions and the subsequently obtained colloidal solution. The characterization of ZnO and MgO PTNPs synthesized from *Moringa oleifera* was carried out by scanning electron microscopy (SEM), dynamic light scattering (DLS) and FT-IR spectroscopy.

FT-IR and SEM analysis

The FT-IR spectrum of the ZnO PTNPs is shown in Fig. 1a. Molecules such as alcohols, phenols, and carboxylic acids present in the plant extract can interact with the zinc surface and contribute to the stabilization of ZnO PTNPs (Shaikhaldein *et al.*, 2021; Fakhari *et al.*, 2019). Characteristic IR bands appeared at 3390 and 2825 cm^{-1} corresponding to the stretching vibra-




Figura 1. Caracterización de fitonanopartículas de ZnO sintetizadas por *M. oleifera*. a) Espectrofotómetro infrarrojo (FT-IR), b) Micrografía electrónica de barrido (SEM), c) Histograma de distribución de tamaño de partícula (DLS).

Figure 1. Characterization of ZnO phytonanoparticles synthesized by *M. oleifera*. a) Infrared spectrum (FT-IR), b) Scanning electron micrograph (SEM), c) Particle size distribution histogram (DLS).

tions of O-H and C-H bonds respectively (Senthilkumar *et al.*, 2017). The band observed at 1500 cm^{-1} can be attributed to the asymmetric and symmetric stretching of carbonyls (C=O) (Moghaddam *et al.*, 2017), while the band at 871 cm^{-1} could be related to the stretching vibrations of Zn-O bonds (Alamdari *et al.*, 2020; Fakhari *et al.*, 2019). On the other hand, the study of MgO PTNPs (Fig. 2a) shows the presence of a broad band of the hydroxyl group at 3381 cm^{-1} (Khajuria *et al.*, 2021; Alamdari *et al.*, 2020). Two distinct bands are observed in the wave number regions at 1429 and 1106 cm^{-1} which are attributed to the bending vibrations of the adsorbed water molecule and the surface hydroxyl group (-OH) (Boro *et al.*, 2020). The band observed at 531 cm^{-1} indicates the formation of the Mg-O bond (Lucho *et al.*, 2018).

SEM imaging of the ZnO PTNPs revealed agglomerated rod-like PTNPs with a diameter range of 3 - 8 nm (Fig. 1b). Velázquez-Gamboa *et al.* (2021) reported the synthesis of ZnO PTNPs from *Moringa oleifera* in the form of rod-shaped, elongated with an average size of 13 nm. The image corresponding to the MgO PTNPs shows that they are irregularly shaped, and highly agglomerated, with an average size of 45 nm (Fig. 2b). Similar results were reported by Amrulloh *et al.* (2021), who synthesized MgO PTNPs from *M. oleifera* with spherical shape and average size ranging from 20 to 80 nm. The study by Gupta *et al.* (2020) highlights the importance of various physicochemical factors in controlling the synthesis of green nanoparticles. In fact, the mentioned factors, including the optimal concentration of metal ions, pH, temperature, concentration of precursors and reducing agents, and reaction time, play fundamental roles in the synthesis of PTNPs, and significantly affect their physicochemical properties and functionality. However, to date, the synthesis and application of specific forms of PTNPs have not been widely reported

or established in the scientific literature. The synthesis and application of nanoparticles, especially in the context of green or environmentally friendly methods, are active areas of research and new developments occur periodically.

DLS and ζ potential analysis

The DLS technique is widely used to determine the hydrodynamic diameter of suspended particles. The average hydrodynamic size of the ZnO PTNPs (Fig. 1c) synthesized in this study was in the range of 9 nm to 30 nm in a stabilized medium. The ζ potential defines the colloidal stability, it is a typical measure of the surface charge of a particle (Mehta *et al.*, 2021). In this study, the ζ potential of ZnO PTNPs in distilled water was +200 mV, therefore, they can be considered as strongly cationic which favors their dispersion in solution. When the particle is partially dispersed in the medium, the ζ potential tends to be positive (Campbell *et al.*, 2019). The exact ζ potential required for stability depends on the specific colloidal system, including the nature of the particles, their size, surface charge, and the surrounding medium. In addition, factors such as temperature and ionic strength can also affect the ζ potential and thus the stability of colloids.

The histogram in Fig. 2c, shows the size distribution of MgO PTNPs, which were observed in the range of 5 - 10 nm with a mean size of 7.5 nm. Moorthy *et al.* (2015) reported the synthesis of MgO PTNPs from *Neem* leaf with a size of 43 nm, while Ali *et al.* (2020) reported the size MgO PTNPs from *Persimmon* extract with a size of 23.88 nm. Regarding the surface charge of the biosynthesized MgO PTNPs MgO a ζ potential of +39.5 mV was determined. These results were similar to those reported by Suresh *et al.* (2014), who reported a ζ potential of +39 mV in MgO PTNPs from *Calotropis gigantean*, indicating the formation of stable nanoparticles in solutions at slightly acidic pH.




Figura 2. Caracterización de fitonanopartículas de MgO sintetizadas por *M. oleifera*. a) Espectro de infrarrojo (FT-IR), b) Micrografía electrónica de barrido (SEM), c) Histograma de distribución de tamaños de partícula (DLS).

Figure 2. Characterization of MgO phytonanoparticles synthesized by *M. oleifera*. a) Infrared spectrum (FT-IR), b) Scanning electron micrograph (SEM), c) Particle size distribution histogram (DLS).

Rebaudioside A, D, E, and M content in elicited *Stevia rebaudiana* Bertoni plants

Recently, some researchers have reported that PTNPs influence the production of secondary metabolites in plants with medicinal properties (Selvakesavan *et al.*, 2023). Rebaudioside A is the most abundant compound in *Stevia rebaudiana* extract; therefore, the commercial value of this product is mainly defined by its content. However, there are other rebaudiosides with similar sweetening power to Reb A, such as Reb D and Reb M, which, although found in low concentrations, could have a significant impact on the sensory properties of *S. rebaudiana*. Regarding the content of rebaudioside A, in Table 1 shows that there is a significant statistical difference ($p < 0.05$) between treatments, where the plants that were elicited with MeJa showed an increase of 18 % with respect to the control plants. These results were consistent with those described by Lucho *et al.* (2018), who reported that the application of this phytohormone induced an increase in rebaudioside A, after 24 and 72 hours of application.

Regarding the treatments with added PTNPs, it was observed that at low concentrations (T3 and T6) there was a significant increase between 4 and 5 %, respectively, compared to the control, while at higher concentrations their content decreased. Javed *et al.* (2017a) reported similar results in the content of steviol glycosides using 1 mg/L ZnO PTNPs in the culture of micropropagated *S. rebaudiana* shoots. These results suggest that PTNPs of both metals at concentrations of 1 mg/L could generate a signaling cascade that probably induces a higher expression or activity of the enzyme UGT76G1. On the other hand, the increase in PTNPs

concentrations could induce greater stress in plants by modifying the metabolism, towards the synthesis of molecules with antioxidant capacity to regulate cell damage (Table 2), thereby inhibiting the synthesis of steviol glycosides.

The results for Reb D and Reb E content were reported as Reb D equivalents, because both molecules are isomers and have the same retention time in the chromatographic analysis used. Table 1 shows that there is no significant statistical difference ($p < 0.05$) between the controls, although the same enzyme catalyzes the glycosylation for the synthesis of these molecules using Reb A as precursor. On the other hand, the application of ZnO and MgO PTNPs, at all the concentrations studied, significantly reduced their content up to 26 %.

As for Reb M, the plants treated with ZnO at 1 mg/mL increased its content by 3 % with respect to the control, while the other treatments did not show significant statistical differences with respect to the control plants.

Several authors have reported that the use of elicitors and phytohormones such as salicylic acid, paclobutrazol, MeJa, some auxins such as gibberellins and nanoparticles (Ag, CuO, ZnO-PEG, ZnO-PVP, CuO-PEG and CuO-PVP) (Gupta *et al.*, 2015; Karimi *et al.*, 2015; Javed *et al.*, 2017a; Javed *et al.*, 2017b; Saraiva *et al.*, 2020; Wang *et al.*, 2021) of chemical, biological and commercial synthesis, affect the synthesis of secondary metabolites, mainly in the production of SGs and compounds with antioxidant capacity. Therefore, ZnO and MgO PTNPs may exhibit similar catalytic properties in certain chemical reactions and could play a similar role in the synthesis of steviol glycosides, depending on the specific reaction conditions and mechanisms involved. However, it is important to note that the catalytic behavior of nanoparticles can

Cuadro 1. Contenido de rebaudiósidos A, D y M en extracto acuoso de hojas secas de *Stevia rebaudiana* Bertoni estimulados con fitonanopartículas de óxido de zinc y magnesio.

Table 1. Rebaudiosides A, D and M content in aqueous extract of *Stevia rebaudiana* Bertoni dried leaves elicited with zinc and magnesium oxide phytonanoparticles.

Treatments	PTNPs mg/L	Rebaudioside A	Rebaudioside D	Rebaudioside M
		mg/ g DW		
T1 Deionized water	0	43.38 c	15.64 a	2.55 b
T2 Metil jasmonate	0	51.19 a	15.82 a	2.44 c
T3 ZnO	1	45.22 b	13.12 b	2.63 a
T4	20	38.8 de	12.05 cd	2.54 b
T5	30	37.67 e	11.71 d	2.51 bc
T6 MgO	1	45.86 b	12.77 bc	2.5 bc
T7	20	42.3 c	12.32 cd	2.52 bc
T8	30	40.49 e	12.4 bcd	2.55 b
	LSD (0.05)	1.73	0.72	0.085

LSD: Least significant difference. Values with the same letter in each column have no statistically significant difference between treatments. T2 (22.43 mg/L Metil jasmonate)

Table 2. Content of gallic acid, chlorogenic acid, caffeic acid and caffeine, in aqueous extract of *Stevia rebaudiana* Bertoni dried leaves elicited with zinc and magnesium oxide phytonanoparticles.

Cuadro 2. Contenido de ácido gálico, ácido clorogénico, ácido cafeico y cafeína en extracto acuoso de hojas secas de *Stevia rebaudiana* Bertoni estimulados con fitonanopartículas de óxido de zinc y magnesio.

Treatments	PTNPs mg/L	Gallig acid	Chlorogenic acid	Caffeic acid	Caffeine
		mg/ kg DW			
T1 Deionized water	0	63.1 a	749.2 de	20.1cd	7.8 b
T2 Metil jasmonate	0	62.3 ab	818.0 d	23 cd	6.0 c
T3 ZnO	1	55.3 c	746.7 e	27.3 c	7.4 bc
T4	20	55.2 c	1777.0 a	26.4 c	10.4 a
T5	30	58.7 bc	1425.2 b	41.6 b	7.4 bc
T6 MgO	1	62.2 ab	142.0 g	17.7 d	5.9 c
T7	20	64.5 a	451.9 f	26.8 c	4.2 d
T8	30	59.2 b	1011.9 c	57.4 a	6.5 bc
	LSD (0.05)	4.82	69.3	1.54	1.54

LSD: Least significant difference. Values with the same letter in each column have no statistically significant difference between treatments. T2 (22.43 mg/L Metil jasmonate)

be highly dependent on several factors, including size, morphology, surface properties, and specific reaction conditions. Lucho *et al.* (2018) reported that MeJa does not negatively affect the expression of genes involved in SGs synthesis. They demonstrated the positive influence of MeJa on the activation of the genes responsible for encoding enzymes in the SGs biosynthesis pathway, reporting that about 12 (80 %) genes catalogued by this elicitor as a strong candidate for use in SGs production optimization. A statistically significant difference was observed in Reb M content, and with these results, it could be assumed that PTNPs act similarly in gene activation; Therefore, PTNPs could induce a higher gene expression of enzymes responsible for stevioside glycosylation, these enzymes are known as UDP-glycosyltransferases.

Gallic acid, chlorogenic acid, caffeic acid, caffeine content

The production of non-enzymatic antioxidants (phenolic compounds) is the protective mechanism against reactive oxygen species (ROS) in plants and their accumulation varies in different environments. Several studies have reported that the use of metal PTNPs induces the synthesis of these metabolites (Gowri *et al.*, 2019; Moharramnejad *et al.*, 2019; Velázquez-Gamboa *et al.*, 2021). Several researchers have reported that metal PTNPs influence the synthesis of secondary metabolites in plants. *Stevia rebaudiana* has an added value because it is a plant that has a high content of phenolic compounds. However, there are some of them with greater antioxidant capacity, such as flavonoids, which, although found in lower concentrations, have a significant impact on the commercial value of the plant.

Near the flavonoid content, those of gallic acid, chlorogenic acid, caffeic acid and caffeine present in the leaves of *S. rebaudiana* nanoelicited are shown in Table 2. It was observed that the control with MeJa did not modify the content of gallic acid, chlorogenic acid, or caffeic acid, only caffeine decreased significantly. However, the elicitation with PTNPs mainly with ZnO at 20 mg/L induced the production of molecules with antioxidant capacity. Chlorogenic acid increased up to 137 %, while caffeic acid increased by 185 %. Likewise, in the results of the caffeine studies, a significant statistical difference was observed between treatment T4 (20 mg/L ZnO PTNPs) and T1 (negative control) showing a 133 % increase. In this case, the results obtained indicate that the application of ZnO and MgO PTNPs induces an increase in antioxidant compounds content. In general, Chandra *et al.* (2017) suggests that PTNPs can stimulate a strong immune response in *Camellia chinensis* plants, leading to the upregulation of defense-related genes, increased activity of defense enzymes, and accumulation of phenolic compounds, all of which play a crucial role in the plant's ability to defend itself against potential threats such as pathogens.

The results of the study indicate that an increase in PTNPs concentration leads to an increase in the production of reactive oxygen species (ROS), but does not reach toxic levels for the plant. Instead, the plant responds by increasing the production of antioxidant compounds. This response

is consistent with the concept of homeostasis in plants. Hussain *et al.* (2021) indicate that oxidative stress was ameliorated in *Persicaria hydropiper* L. seedlings exposed to ZnO NPs through increased production of free proline, phenolics, flavonoids, and the activation of antioxidant enzymes.

CONCLUSIONS

The elicitation of *S. rebaudiana* with ZnO and MgO PTNPs induces the biosynthesis of rebaudioside A and M, which could improve the sweetening potential of its extracts, in addition to a significant increase in the content of chlorogenic acid and caffeic acid, molecules that contribute to the antioxidant activity. Consequently, the application of ZnO and MgO PTNPs to *S. rebaudiana* plants probably gives added commercial value to the crop. However, it is necessary to continue the studies to evaluate the phytotoxic effect of the extract obtained from plants treated with these PTNPs.

ACKNOWLEDGMENTS

The authors thank the Instituto Tecnológico de Tuxtla Gutierrez for providing the facilities

FUNDING

The authors thank CONAHCyT Mexico for providing financial support to Velázquez-Gamboa with Scholarship: 849710 to carry out this research.

CONFLICTS OF INTEREST

Consent for Publication: All authors agreed to publish this paper in this journal.

Ethical approval: This study does not contain any studies with human participants or animals performed.

Conflict of interest: The authors declare no conflict of interest.

REFERENCES

- Abdel-Aziz, M.M., Emam, T.M., and Elsherbiny, E.A. 2020. Bioactivity of magnesium oxide nanoparticles synthesized from cell filtrate of endobacterium *Burkholderia rinojensis* against *Fusarium oxysporum*. *Materials Science and Engineering C*, 109, 110617. <https://doi.org/10.1016/j.msec.2019.110617>
- Alamdari, S., Ghamsari, M.S., Lee, C., Han, W., Park, H.H., Tafreshi, M.J., Afarideh, H. and Ara, M.H. 2020. Preparation and characterization of zinc oxide nanoparticles using leaf extract of *Sambucus ebulus*. *Applied Sciences*, 10(10), 3620. <https://doi.org/10.3390/app10103620>
- Ali, R., Shanan, Z.J., Saleh, G.M. and Abass, Q. 2020. Green synthesis and the study of some physical properties of MgO nanoparticles and their antibacterial activity. *Iraqi Journal of Science*, 61(2), 266-276. <https://doi.org/10.24996/ij.s.2020.61.2.9>
- Amrulloh, H., Fatiqin, A., Simanjuntak, W., Afriyani, H. and Annisa, A. 2021. Antioxidant and antibacterial activities of magnesium oxide nanoparticles prepared using aqueous extract of *Moringa oleifera* Bark as green agents. *Journal of Multidisciplinary Applied Natural Science*, 1(1), 44-53. <https://doi.org/10.47352/jmans.v1i1.9>



- Boro, B., Nath, A.K., Barthakur, M. and Kalita, P. 2020. Synthesis and characterization of MgO nanoparticle and its in vitro cytotoxic effect on erythrocytes. In: Ramkrishna, D., Sengupta, S., Dey Bandyopadhyay, S., Ghosh, A. (eds) *Advances in Bioprocess Engineering and Technology. Lecture Notes in Bioengineering*. Springer, Singapore. pp. 199-207. https://doi.org/10.1007/978-981-15-7409-2_20
- Bouttier-Figueroa, D.C., Cortez-Valadez, M., Flores-Acosta, M. and Robles-Zepeda, R.E. 2023. Synthesis of metallic nanoparticles using plant's natural extracts: Synthesis mechanisms and applications. *Biotecnia*, 25(3): 125-139. <https://doi.org/10.18633/biotecnia.v25i3.1916>
- Campbell, J., Burkitt, S., Dong, N. and Zavaleta, C. 2019. Nanoparticles for biomedical applications, chapter 9: nanoparticle characterization techniques (Eun, J. C., Lorraine, L. Carlos R., ed.), Elsevier. pp. 129-144. <https://doi.org/10.1016/B978-0-12-816662-8.00009-6>
- Cauch-Cauch, R., Tun-Suárez, J.M., Cristóbal-Alejo, J., Herrera-Parra, E., Andueza-Noh, R. and Lozano-Contreras, M.G. 2022. Evaluación de consorcios micorrízicos arbusculares nativos en interacción con niveles de fósforo en la promoción del crecimiento y fotosíntesis de *Stevia rebaudiana* Bertoni. *Biotecnia* 25(1): 67-68. <https://doi.org/10.18633/biotecnia.v25i1.1765>
- Chandra, S., Chakraborty, N., Panda, K. and Acharya, K. 2017. Chitosan-induced immunity in *Camellia sinensis* (L.) O. Kuntze against blister blight disease is mediated by nitric-oxide. *Plant Physiology and Biochemistry*, 115, 298-307. <https://doi.org/10.1016/j.plaphy.2017.04.008>
- Chaudhuri, S.K. and Malodia, L. 2017. Biosynthesis of zinc oxide nanoparticles using leaf extract of *Calotropis gigantea*: Characterization and its evaluation on tree seedling growth in nursery stage. *Applied Nanoscience*, 7, 501-512. <https://doi.org/10.1007/s13204-017-0586-7>
- Fakhari, S., Jamzad, M. and Kabiri, F.H. 2019. Green synthesis of zinc oxide nanoparticles: a comparison. *Green Chemistry Letters and Reviews*, 12, 19-24. <https://doi.org/10.1080/17518253.2018.1547925>
- Gasmalla, M.A., Yang, R., Musa, A., Hua, X. and Ye, F. 2017. Influence of sonication process parameters to the state of liquid concentration of extracted rebaudioside A from *Stevia rebaudiana* Bertoni leaves. *Arabian Journal of Chemistry*, 10(5), 726-731. <https://doi.org/10.1016/j.arabjc.2014.06.012>
- Gowri, M., Latha, N. and Rajan, M. 2019. Copper oxide nanoparticles synthesized using *Eupatorium odoratum*, *Acanthospermum hispidum* leaf extracts, and its antibacterial effects against pathogens: A comparative study. *BioNanoScience*, 9, 545-552. <https://doi.org/10.1007/s12668-019-00655-7>
- Gupta, E., Purwar, S., Sundaram, S. and Rai, G.K. 2013. Nutritional and therapeutic values of *Stevia rebaudiana*: A review. *Journal of Medicinal Plants Research*, 7(46), 3343-3353. <https://doi.org/10.5897/JMPR2013.5276>
- Gupta, P., Sharma, S. and Saxena, S. 2015. Biomass yield and steviol glycoside production in callus and suspension culture of *Stevia rebaudiana* treated with proline and polyethylene glycol. *Applied Biochemistry and Biotechnology*, 176, 863-874. <https://doi.org/10.1007/s12010-015-1616-0>
- Gupta, M., Tomar, R.S., and Mishra, R.K. 2020. Factors affecting biosynthesis of green nanoparticles. *Our Heritage*, 68(30), 10530-10555.
- Hussain, F., Hadi, F. and Rongliang, Q. 2021. Effects of zinc oxide nanoparticles on antioxidants, chlorophyll contents, and proline in *Persicaria hydropiper* L. and its potential for Pb phytoremediation. *Environmental Science and Pollution Research*, 28, 34697-34713. <https://doi.org/10.1007/s11356-021-13132-0>
- Javed, R., Usman, M., Yücesan, B., Zia, M. and Gürel, E. 2017a. Effect of zinc oxide (ZnO) nanoparticles on physiology and steviol glycosides production in micropropagated shoots of *Stevia rebaudiana* Bertoni. *Plant Physiology and Biochemistry*, 110, 94-99. <https://doi.org/10.1016/j.plaphy.2016.05.032>
- Javed, R., Zia, M., Yücesan, B. and Gürel, E. 2017b. Abiotic stress of ZnO-PEG, ZnO-PVP, CuO-PEG and CuO-PVP nanoparticles enhance growth, sweetener compounds and antioxidant activities in shoots of *Stevia rebaudiana* Bertoni. *IET Nanobiotechnology*. 11, 898-902. <https://doi.org/10.1049/iet-nbt.2016.0247>
- Karimi, M., Hashemi, J., Ahmadi, A., Abbasi, A., Pompeiano, A., Tavarini, S., Guglielminetti, L. and Angelini, L.G. 2015. Opposing effects of external gibberellin and daminozide on *Stevia* growth and metabolites. *Applied Biochemistry and Biotechnology*, 175, 780-791. <https://doi.org/10.1007/s12010-014-1310-7>
- Khajuria, A.K., Kumari, M., Kandwal, A., Singh, A. and Bisht, N.S. 2021. Biofabrication of zinc oxide nanoparticles from two different zinc sources and their antimicrobial activity. *BioNanoScience*. 11, 793-809. <https://doi.org/10.1007/s12668-020-00802-5>
- Krishnani, K.K., Boddu, V.M. and Chadha, N.K. 2022. Metallic and non-metallic nanoparticles from plant, animal, and fisheries wastes: potential and valorization for application in agriculture. *Environmental Science and Pollution Research*, 29, 81130-81165. <https://doi.org/10.1007/s11356-022-23301-4>
- Lucho, S.R., Do-Amaral, M.N., Milech, C., Ferrer, M.Á., Calderón, A.A., Bianchi, V.J. and Braga, J.B. 2018. Elicitor-induced transcriptional changes of genes of the steviol glycoside biosynthesis pathway in *Stevia rebaudiana* Bertoni. *Journal of Plant Growth Regulation*, 37, 971-985. <https://doi.org/10.1007/s00344-018-9795-x>
- Magdaleno García, G., Juárez Maldonado, A., Betancourt Galindo, R., González Morales, S., Cabrera de la Fuente, M., Sánchez Vega, M. and Méndez, A. 2023. Zinc oxide nanoparticle morphology modify germination and early growth of bell pepper seedlings. *Biotecnia* 25(3): 5-1. <https://doi.org/10.18633/biotecnia.v25i3.1908>
- Masarovicová, E. and Králová, K. 2013. Metal nanoparticles and plants. *Ecological Chemistry and Engineering S*, 20(1), 9-22. <https://doi.org/10.2478/eces-2013-0001>
- Mehta, M.R., Mahajan, H.P. and Hivrale, A.U. 2021. Green synthesis of chitosan capped-copper nano biocomposites: synthesis, characterization, and biological activity against plant pathogens. *BioNanoScience*, 11, 417-427. <https://doi.org/10.1007/s12668-021-00823-8>
- Moghaddam, A.B., Moniri, M., Azizi, S., Rahim, R.A., Ariff, A., bin-Saad, W.Z., Namvar, F., Navaderi, M. and Mohamad, R. 2017. Biosynthesis of ZnO nanoparticles by a new *Pichia kudriavzevii* yeast strain and evaluation of their antimicrobial and antioxidant activities. *Molecules*. 22(6), 872. <https://doi.org/10.3390/molecules22060872>

- Moharramnejad, S., Azam, A.T., Panahandeh, J., Dehghanian, Z. and Ashraf, M. 2019. Effect of methyl jasmonate and salicylic acid on in vitro growth, stevioside production, and oxidative defense system in *Stevia rebaudiana*. Sugar Tech, 21, 1031-1038. <https://doi.org/10.1007/s12355-019-00727-8>
- Moorthy, S.K., Ashok, C.H., Rao, K.V. and Viswanathan, C. 2015. Synthesis and characterization of MgO nanoparticles by neem leaves through green method. Mater Today, 2, 4360-4368. <https://doi.org/10.1016/j.matpr.2015.10.027>
- Nongbet, A., Mishra, A.K., Mohanta, Y.K., Mahanta, S., Ray, M.K., Khan, M., Baek, K.H. and Chakrabartty, I. 2022. Nanofertilizers: A smart and sustainable attribute to modern agriculture. Plants, 11(19), 2587. <https://doi.org/10.3390/plants11192587>
- Stevia Market Outlook from 2024 to 2034. [Consultado 06 de agosto 2024] 2024. <https://www.futuremarketinsights.com/reports/global-stevia-market>
- Pandey, D.K. and Dey, A. 2018. HPTLC Analysis of the antioxidant and possible antidiabetic chlorogenic acid in the in situ and in vitro populations of the low-calorie sweetener *Stevia rebaudiana* (Bert.) Bertoni. Analytical Chemistry Letters, 8, 872-881. <https://doi.org/10.1080/22297928.2018.1515657>
- Ramezani, M., Asghari, S., Gerami, M., Ramezani, F. and Karimi, A.M. 2020. Effect of silver nanoparticle treatment on the expression of key genes involved in glycosides biosynthetic pathway in *Stevia rebaudiana* B. Plant Sugar Tech, 22, 518-527. <https://doi.org/10.1007/s12355-019-00786-x>
- Raliya, R., Tarafdar, J.C., Choudhary, K., Mal, P., Raturi, A., Gautam, R. and Singh, S.K. 2014. Synthesis of MgO nanoparticles using *Aspergillus tubingensis* TFR-3. Journal of Bionanoscience, 8, 34-38. <https://doi.org/10.1166/jbns.2014.1195>
- Rodríguez-Pérez, C., Quirantes-Piné, R., Fernández-Gutiérrez, A. and Segura-Carretero, A. 2015. Optimization of extraction method to obtain a phenolic compounds-rich extract from *Moringa oleifera* Lam leaves. Industrial Crops and Products, 66, 246-254. <https://doi.org/10.1016/j.indcrop.2015.01.002>
- Saraiva, A., Carrascosa, C., Raheem, D., Ramos, F. and Raposo, A. 2020. Natural sweeteners: The relevance of food naturalness for consumers, food security aspects, sustainability and health impacts. International Journal of Environmental Research and Public Health, 17(17):6285. <https://doi.org/10.3390/ijerph17176285>
- Selvakesavan, R.K., Kruszka, D., Shakya, P., Mondal, D. and Franklin, G. 2023. In nanomaterial interactions with plant cellular mechanisms and macromolecules and agricultural implications, chapters 2: impact of nanomaterials on plant secondary metabolism. (Jameel, M. Lina, M. Mohan, J. ed), Springer Cham pp. 133-170. https://doi.org/10.1007/978-3-031-20878-2_6
- Senthilkumar, N., Nandhakumar, E., Priya, P., Soni, D., Vimalan, M. and Vetha, I. 2017. Synthesis of ZnO nanoparticles using leaf extract of: *Tectona grandis* (L.) and their anti-bacterial, anti-arthritis, antioxidant and in vitro cytotoxicity activities. New Journal of Chemistry, 41, 10347-10356. <https://doi.org/10.1039/c7nj02664a>
- Shaikhaldein, H.O., Al-qurainy, F., Khan, S., Nadeem, M., Tarroum, M., Salih, A.M., Gaafar, A.Z., Alshameri, A., Alansi, S., Alenezi, N.A. and Alfarraj, N.S. 2021. Biosynthesis and characterization of ZnO nanoparticles using *Ochradenus arabicus* and their effect on growth and antioxidant systems of *Maerua oblongifolia*. Plants, 10(9), 1808. <https://doi.org/10.3390/plants10091808>
- Suresh, J., Yuvakkumar, R., Sundrarajan, M. and Hong, S.I. 2014. Green synthesis of magnesium oxide nanoparticles. Advanced Materials Research, 952, 141-144. <https://doi.org/10.4028/www.scientific.net/AMR.952.141>
- Velázquez-Gamboa, M.C., Rodríguez-Hernández, L., Abud-Archila, M., Gutiérrez-Miceli, F.A., González-Mendoza, D., Valdez-Salas, B., González-Terreros, E. and Luján-Hidalgo, M.C. 2021. Agronomic biofortification of *Stevia rebaudiana* with zinc oxide (ZnO) phytonanoparticles and antioxidant compounds. Sugar Tech, 23, 453-460. <https://doi.org/10.1007/s12355-020-00897-w>
- Vijayakumar, S., Punitha, V.N. and Parameswari, N. 2022. Phytonanosynthesis of MgO nanoparticles: Green Synthesis, characterization and antimicrobial evaluation. Arabian Journal for Science and Engineering, 47, 6729-6734. <https://doi.org/10.1007/s13369-021-06107-3>
- Wang, J., Li, J., Li, J., Li, J., Liu, S., Huang, L. and Gao, W. 2017. Production of active compounds in medicinal plants: from plant tissue culture to biosynthesis. Chinese Herbal Medicines, 9(2), 115-125. [https://doi.org/10.1016/s1674-6384\(17\)60085-6](https://doi.org/10.1016/s1674-6384(17)60085-6)
- Wang, Y., Sun, X., Jia, X., Zhu, L. and Yin, H. 2021. Comparative transcriptomic of *Stevia rebaudiana* provides insight into rebaudioside D and rebaudioside M biosynthesis. Plant Physiology and Biochemistry, 167, 541-549. <https://doi.org/10.1016/j.plaphy.2021.08.028>