

Response to bocashi and the vermicompost of *Moringa oleifera* Lam. after pruning

Iván Francisco Mota Fernández¹
Ofelia Andrea Valdés Rodríguez¹
Gabina Sol Quintas²
Arturo Pérez Vázquez^{3§}

¹The College of Veracruz. Carrillo Port no. 26, Central Zone, Xalapa Enríquez, Veracruz, Mexico. CP. 91000. Tel. 01 (228) 1490182. (ivan.mota.im@gmail.com, andrea.valdes@gmail.com). ²Technological Institute of Perote. Perote, Veracruz. Tel. 01 (228) 8264650. (macadamiamexico@yahoo.com.mx). ³Postgraduate School-*Campus* Veracruz. Road Veracruz-Xalapa km 13.5, Veracruz, Mexico.

§Corresponding author: parturo@colpos.mx.

Abstract

Moringa oleifera Lam. due to its multiple uses, it has an enormous potential as an alternative for sustainable cultivation to be implemented in the central area of the state of Veracruz. The present work evaluates the vegetative and reproductive response of the species to organic fertilization after pruning in this region. The experimental design was completely randomized with three-year-old moringa trees, pruned only once at a height of 1.5 m, to which two organic fertilizers were applied: vermicompost and bocashi and one control. The fertilizers were applied at the beginning of the experiment and later every four months. The variables: 1) number of sprouts; 2) survival of sprouts; 3) length of sprouts; 4) thickness of sprouts; 5) number of bunches per plant; 6) number of buttons per cluster; 7) flowers per bunch; 8) fruits per bunch; 9) number of fruits per plant; 10) length of fruits; 11) number of seeds per fruit; and 12) seed dimension, were measured every month for 10 months. Treatments were evaluated by one-way analysis of variance ($p < 0.05$). The results showed that the application of organic fertilizers did not influence variables 3, 4, 5, 6, 7 and 8 as there were no significant statistical differences in these; except the total number of fruits per plant, where plants with bocashi achieved the highest productivity. It is concluded that the best fertilizer for fruit production was bocashi, while in vermicompost no significant differences were observed.

Keywords: *Moringa oleifera* Lam., organic fertilizers, vegetative response.

Reception date: January 2019

Acceptance date: March 2019

Introduction

Moringa (*Moringa oleifera* Lam.) is a tree native to northwestern India (Zayed, 2012), arrived in Mexico probably by boats made by Filipino sailors traveling from Manila, Philippines to Acapulco, Guerrero (Olson and Fahey, 2011). Moringa is very appreciated for its nutritional and medicinal characteristics, present in most of the parts that make it up (leaves, flowers and fruits, all edible). The versatility of environmental conditions to which moringa plant adapts, including the tropical zones of low humidity in the soil, high temperatures, high evaporation and large variations in precipitation (Alves *et al.*, 2010), turn it into a species with great potential to grow in different areas of our country.

Although this plant, in particular, develops better under 500 masl and not above 1 500 masl (Olson and Fahey, 2011). Some authors indicate that Moringa requires a minimum of 250 mm of annual rainfall, and can tolerate clay and sandy soils (Valdes-Rodríguez *et al.*, 2014), but poorly drained soils impair its growth (Velázquez *et al.*, 2016). With respect to their nutritional requirements, there are studies related to organic fertilization in moringa, which show their positive impact on the production of foliar biomass mainly for human and animal consumption (cattle) (Pérez *et al.*, 2010; Suárez, 2014).

Organic fertilization is important if you want to establish a sustainable crop or with low environmental impacts. An organic fertilizer originates through the aerobic and anaerobic decomposition of organic waste mainly of animal and vegetable origin. This action occurs through microorganisms that exist in the waste itself, under controlled conditions, which produces a partially stable material of slow decomposition under favorable conditions (Restrepo, 2001).

Two organic fertilizers widely known in Mexico and much of Latin America are the bocashi and the vermicompost, their simple way of processing them and low production cost, make them a viable option for farmers who use them. The word bocashi comes from the Japanese language and, in the case of fermented organic fertilizers, means to pre-steam the organic materials of the fertilizer through the heat generated by the decomposition (Restrepo and Hensel, 2009). Likewise, the vermicompost is a product resulting from the microbial transformation of organic waste in controlled environments and is obtained when several species of earthworms participate in its elaboration (Olivares *et al.*, 2012).

The positive impact of organic fertilizers in degraded environments is an alternative that is important to promote and increase, since with its application in the soil the growth of the plants is stimulated, the fertility, nutrition and the structure of the same are gradually improved, as well the water retention capacity and the activity of microorganisms are increased, which contributes to reduce the loss of nutrients by leaching (Restrepo and Hensel, 2009). Therefore, it is necessary to promote the application of organic fertilizers in places and crops with productive potential, as is the case of moringa, and thus avoid the application of agrochemicals, which are one of the main causes of degradation and pollution of natural resources and biodiversity (Altieri, 1999; Ramos and Terry, 2014).

Currently there are few studies on moringa reported in the state of Veracruz. One of them was performed by Valdes *et al.* (2014) and is focused on the registration of moringa allometric measurements (growth, flowering and fruit production) without receiving organic fertilization. Another work done by Quintas *et al.* (2016) analyzed the vegetative response of moringa to two organic fertilizers. However, this only comprised a period of seven months and did not address the response of such fertilizers on the reproductive variables. Therefore, an evaluation of the productive variables would allow estimating the effects of organic fertilizers on the productivity of moringa plants, in order to determine if their application represents an efficient and profitable agronomic practice for this crop.

The objective of this research was to evaluate the use of organic fertilizers (vermicompost and bocashi) on the vegetative and reproductive response of moringa after pruning, in the central region of the state of Veracruz.

Materials and methods

Study area

This research was conducted at the Postgraduate College (COLPOS), Campus Veracruz, which is located in the municipality of Manlio Fabio Altamirano, in the central area of the state of Veracruz. This municipality is located at the coordinates 19°18' North latitude and 96° 32' West longitude. It has an altitude of 25 m and its soil type is loam-clay-sandy. The climate is warm with rains in summer and winter; the average annual rainfall of 1 239.5 mm and the average annual temperature of 25 °C (Cruz *et al.*, 2015).

Experimental procedure

Three-year-old moringa plants were evaluated, planted at a density of 1 m by 1 m, with an average height of 6 m. 24 individuals were randomly selected to apply the treatments, with eight individuals per block. The first treatment consisted of the application of vermicompost made with 30% coffee pulp and 70% cane bagasse. The second treatment consisted in the application of bocashi, made with cattle manure (50 kg), maize stubble (25 kg), litter (25 kg), ash (10 kg), wheat bran (25 kg), molasses (4 L), bread yeast (250 gr) and soil of the region (50 kg) and finally the control, to which no fertilizer was applied.

In order to standardize the size of the plants and facilitate the taking of measurements, a single pruning was performed one day before starting the treatment, the trees being in a period of vegetative growth without evidence of floral clusters and fruits. The pruning consisted in the cutting of the main trunk at a height of 1.5 m and the elimination of the lateral branches and sprouts of the plant, leaving only the main trunk. The next day the first application of organic fertilizers was made. Three applications of four kilograms per plant were made every four months distributed during the ten months that the work lasted.

The fertilizer was applied in a radius of 40 cm of the main trunk of the plant, it was buried 10 cm and later it was covered with the soil removed from said area. After the application of the organic fertilizer a single irrigation was practiced using 19 L of water to irrigate each plant. Between the application periods no irrigation was done and the weed control was done manually.

Additionally, soil sampling of the experimental site was carried out and the chemical analysis of the organic fertilizers involved was carried out. For the soil, 15 sub-samples of 500 g were randomly collected, each one at a depth of 10-15 cm from the soil, the samples were combined in a homogeneous manner and a 500 g portion was subsequently taken. Likewise, for organic fertilizers (vermicompost and bocashi) a representative sample of (500 g) was taken. The soil samples of the region and the organic fertilizers were sent for analysis to the soil laboratory of the Institute of Ecology, AC (INECOL).

The methodology used for the sample of soil and organic fertilizers was the one indicated by NOM-021REC/NAT 2000. The variables evaluated were divided into A) vegetative, which were analyzed weekly for four months: 1) number of sprouts; 2) survival of sprouts; 3) length of sprouts; and 4) regrowth thickness; and B) reproductive: 5) number of bunches per plant; 6) number of buttons per cluster; 7) number of flowers per bunch; and 8) number of fruits per bunch; 9) number of fruits per plant; 10) length of fruits; 11) number of seeds per fruit; and 12) seed dimension. These last variables were evaluated during a period of ten months starting from March to December 2016.

Vegetative variables

The number of sprouts and their survival per plant was determined by means of the total accounting of these on their initial registration and was carried out weekly for four months. The measurement that corresponds to the variables of length and thickness of the sprouts was taken with the help of a flexometer (1 mm precision) and a digital vernier (0.01 mm precision) respectively, for which six sprouts were selected per plant, because this was the minimum number of sprouts surviving per plant the first two months after pruning. During the first five months, these two variables were evaluated weekly; however, when observing that moringa's growth was slow, measurements were made monthly from the sixth month.

Reproductive variables

The total number of floral clusters and fruits per plant was counted weekly. Of the total floral clusters 10 were selected for each plant to be monitored weekly. From each floral cluster selected, the number of buttons, flowers and fruits was counted. Also, the fruits were measured in length, using a flexometer (1 mm) to measure each pod from one end to the other. To determine the total number of seeds per pod all the fruits of the harvest were opened and all the seeds were extracted in order to be counted, later they were measured with a digital vernier (0.01 mm precision) of length and width.

Statistical analysis

For each variable, the corresponding descriptive statistics were obtained (average, standard deviation, maximum, minimum and coefficient of variation). The analysis of each variable between treatments was carried out using a one-way analysis of variance test with a level of significance of 5%. The post-hoc tests were performed using the Tukey method, both analyzes supported by the SigmaPlot 10.0 software.

Results and discussion

The results of the physical-chemical analysis of the soil of the region and the two organic fertilizers showed a remarkable difference in three P, N and Mg macronutrients, with the vermicompost obtaining the highest figures of these. According to González-González and Crespo-López (2016), percentages of N from 1.5% in organic fertilizers favor the production of foliar biomass up to 100%. So the vermicompost turned out to be the fertilizer with the greatest potential for foliage production. While Sarwar *et al.* (2017) established that doses less than 1% of N-P-K can increase moringa stems and foliage by 50% or more in a free soil without fertilization.

So the doses found in both vermicompost and bocashi have the ability to increase the growth of plants above the control. Additionally, it is considered that the pH of bocashi would tend to improve the alkalinity of the soil of the site, which was found to be acidic (Table 1).

Table 1. Physical and chemical composition of the soil of the region and the organic fertilizers used.

Component	Control	Bocashi	Vermicompost
pH	5.63	8.23	5.12
P available (mg kg ⁻¹)	7	26	292
P total (mg kg ⁻¹)	254	1867	2788
N (%)	0.17	0.95	2.79
K (cmol kg ⁻¹)	0.41	18.68	9.76
MO (%)	3.03	17.08	41.51
C/N	12	16	10
Ca (cmol kg ⁻¹)	10.28	34.47	31.28
Mg (cmol kg ⁻¹)	4.93	14.5	18.74
Fe (mg kg ⁻¹)	32.3	149.5	342.8
Mn (mg kg)	23.7	23.6	61.8
Cu (mg kg ⁻¹)	0.31	2.5	6.2
Zn (mg kg ⁻¹)	0.88	17.84	19.87
Carbon organic (%)	1.76	9.91	24.08
Density (g cm ⁻³)	1.51	0.98	0.68
Clay (%)	51.6	21.6	27.6
Slime (%)	26.4	33.6	19.6
Sand (%)	22	44.7	52.7

While the high levels of Fe in bocashi and vermicompost may favor the synthesis of chlorophyll from moringa plants. Likewise, the high contents of Mn will favor the photosynthesis and assimilation of nitrogen (Azcon and Talon, 2013).

Vegetative variables

After pruning and at the end of a period of 16 weeks it was observed that the treatment with bocashi had an average of 12.87 sprouts per plant, vermicompost 8.87 and control 13.12. In this variable, no statistical differences were found between treatments ($p= 0.079$). This result could be due to the trunk diameters of the subjects fertilized with vermicompost were 18% lower than those of the control, while in bocashi they were 15% lower than the control, which is related in the increase to the number of sprouts, which was proportional to the diameter of the stems.

In this regard, Medina *et al.* (2007) obtained an average of 11 to 12 sprouts per plant without the application of any type of fertilizer, which is similar to what was obtained in this experiment. The percentage of surviving sprouts per plant was 73% for bocashi, 66% for vermicompost and 77% for control and no significant statistical differences were found between treatments ($p= 0.867$). No statistically significant differences were found between treatments in the length of sprouts ($p= 0.174$) and thickness of sprouts ($p= 0.514$). While the average number of surviving sprouts per plant until day 283 of monitoring was 3.87, a result lower than that found by Swati *et al.* (2013), which was of eight sprouts with application of organic fertilization.

The results obtained indicate that the low survival of the sprouts could be due to the distance in which the moringa plants were planted, which was only 1 m between each of them, being a factor that could negatively affect this variable, the main cause observed was the fall of the sprouts due to wilting. Previous research mentions that the height of the plants and the number of sprouts that may come to have is related to the distance of planting, being increased if it is greater between plant and plant (Sosa *et al.*, 2016), thus reaching higher yields according to the density in which the plants are planted (Lok and Suárez, 2014). From the previous results it is deduced that the vegetative response to the treatments was not significant in a period of 10 months.

Therefore, if moringa is to be used under similar conditions in a similar period of time, it is not possible to recommend the application of these fertilizers in the quantities applied for a greater production of biomass. This is explained because both vermicompost and bocashi are slow release and require periods of one to three years of constant application to observe results and thus show their effectiveness (Restrepo and Hensel, 2006; Ramos and Terry, 2014), so that with this characteristic they could contribute substantially to the improvement of the biological activity of the soil and therefore of the plant (Usman *et al.*, 2003), but in a period of time of at least two years.

However, the greater average in the variable of length of the regrowths was obtained vermicompost with 13% greater than the control and 18% greater than the bocashi; in addition, in the thickness variable of the regrowth, vermicompost achieved 14% and 16% more than the control and bocashi, respectively (Figure 1 and 2).




Figure 1. Growth of length of sprouts of *M. oleifera* by treatment. The sidebars represent the standard error of the mean.




Figure 2. Growth of the regrowth thickness of *M. oleifera* by treatment. The sidebars represent the standard error of the mean.

This behavior, although not yet significant, could be due to the fact that organic fertilizers indicated a higher percentage of Nitrogen (N) in vermicompost, which was 66% higher than bocashi and 94% higher than control. In this regard, it is known that N is the nutrient with greater importance for the development of the plant, along with P and potassium (K) (Azcon and Talon, 2013).

Reproductive variables

There were no significant statistical differences in the number of floral clusters per plant ($p=0.183$), number of buttons per floral cluster ($p=0.511$), flowers per floral cluster ($p=0.605$) and fruits per floral cluster ($p=0.664$). However, bocashi was the only treatment that

consolidated mature fruits to be harvested (Table 2) and the total number of fruits per plant ($p=0.037$) was the only variable that had significant statistical difference, with the treatment with bocashi giving the greater number.

Table 2. Average values and standard deviation (dev) of the variables evaluated in the different treatments applied to *Moringa oleifera* plants.

Variable	Control		Vermicompost		Bocashi				
	Average	±desv	Average	±desv	Average	±desv			
Length of regrowth	218.34	±96.93	a	253.28	±108	a	207.28	±77.41	a
Thickness of sprouts	27.53	±10.59	a	32.04	±15.1	a	26.84	±10.69	a
Total number of bunches per plant	14.43	±15.47	a	8.73	±9.99	a	14.3	±18.21	a
Buttons by cluster	18.21	±33.85	a	7.71	±19.48	a	16.12	±32.08	a
Flowers per cluster	3.1	±6.99	a	1.61	±6.23	a	2.64	±7.26	a
Fruits per cluster	0.46	±1.4	a	0.61	±1.3	a	0.85	±1.56	a
Total number of fruits per plant	1.89	±3.07	ab	0.94	±2.18	b	3.32	±7.35	a

Different letters in the columns represent statistical differences between treatments ($p < 0.05$).

This result could be due to the fact that bocashi has microorganisms capable of solubilizing certain minerals such as Ca and phosphates, the latter being determinants for flower growth and fruit mooring (Rodríguez *et al.*, 2005).

It is also possible that the difficulty to detect significant differences in the evaluated parameters was due to the enormous variability found in the response of all the plants, with high coefficients of variation (Table 3), where they could influence the climatological conditions and the intrinsic variability of the plants of this origin (Table 3) (Toral *et al.*, 2013).

Table 3. Maximum and minimum values of the variables evaluated in the different treatments applied to *Moringa oleifera* plants.

Variable	Control		Vermicompost		Bocashi	
	Max-min	CV	Max-min	CV	Max-min	CV
Length of regrowth	385-42	0.44	455-90	0.42	397-70	0.37
Thickness of sprouts	52-11	0.38	66-12	0.47	55-7	0.39
Total number of bunches per plant	77-0	1.07	47-0	1.14	100-0	1.27
Buttons by cluster	134-0	1.85	113-0	2.52	156-0	1.99
Flowers per cluster	36-0	2.25	58-0	3.86	55-0	2.75
Fruits per cluster	9-0	3.04	7-0	2.11	10-0	1.84
Total number of fruits per plant	12-0	1.63	9-0	2.33	30-0	2.21

Length of the fruits

Seven months after the pruning, a single harvest was carried out, where 24 fruits were obtained, all from the treatment fertilized with bocashi, since the control and the vermicompost did not present mature fruits.

These results could be due to the higher content of K and Ca, as well as silt and a higher pH present in the bocashi, which facilitated the assimilation of these nutrients strongly related to flower arrest and fruit development (Azcon-Bieto and Talon, 2013). The average length of the moringa pod was 27.39 cm, measured within the parameters reported for this plant, ranging from 10 cm to 50 cm in length (Olson and Fahey, 2011).

Although this length of the pods was 10 cm less than that obtained by other authors who also fertilized with mixtures that included vermicompost, but for a period of one year (Swati *et al.*, 2013). Therefore, it is considered necessary to evaluate organic fertilizers for a period longer than one year in order to determine if there are positive long-term development effects, given the slow release of these biofertilizers. Additionally, it has been documented that moringa develops better in light soils, and in this site the substrate was mostly clayey, which makes the penetration of the roots difficult, since it is a heavy and acid soil, therefore the use of nutrients is made more slower (Pérez *et al.*, 2010).

Number of seeds

The only treatment that managed to consolidate ripe fruit was the fertilization with bocashi, from which a minimum of eight and a maximum of 23 seeds per fruit were obtained, with an average of 14.7 ± 3.3 seeds per fruit. These values were similar to that reported by Pérez *et al.* (2010), ranging from 12 to 25 seeds, and slightly higher than that found by Pérez *et al.* (2010), ranging from 10 to 20 seeds, who describe the typical production of these plants. In a similar study where organic fertilization was also used in moringa plants, a maximum of 14 seeds and a minimum of seven seeds per fruit Swati (2013) was found, from which it is inferred that this bocashi substantially improved the number of seeds produced in the place.

Seed measurement

The average size of the seeds was 10.85 mm long and 10.66 mm wide. Data that coincide with moringa seeds before being sown, which were 12.9 ± 1.1 mm in length and 11.4 ± 0.8 mm in width and 268.9 ± 42.9 mg in weight (Valdes *et al.*, 2014). In another work similar measures (11.34 to 15.16 mm and 9.77 to 12.19 mm) were observed to what was found in this study (Oloyede *et al.*, 2015), so the application of organic fertilizers did not seem to affect the dimensions of the seeds.

Conclusions

Organic bocashi and vermicompost fertilizers applied for ten months in moringa did not show significant effects on the vegetative variables of *Moringa oleifera*. On the other hand, in the reproductive ones, differences were noted in the total number of fruits per plant when fertilized with bocashi. For this reason, the application of bocashi is recommended to improve the production of fruits of the plant. In the same way, when organic fertilizers are used in heavy soils, it is recommended to evaluate the plants for a period greater than a year, to determine if they have better effects on the vegetative variables.

Cited literature

- Altieri, M. 1999. Agroecología: bases científicas para una agricultura sustentable. Agroforestry systems Montevideo: Editorial Nordan-Comunidad. Volumen 35. 338 p.
- Alves, I.; Silva, J.; Carneiro, P. e Bakke, O. 2010. Características de crescimento e valor forrageiro da moringa (*Moringa oleifera* Lam.) submetida a diferentes adubos orgânicos e intervalos de corte. Engenharia Ambiental-Espírito Santo Do Pinhal. 7(2):133-144.
- Azcón, J. y Talón, M. 2013. Fundamentos de fisiología vegetal. McGrawHill. 2^{da} (Ed). Madrid. 522 p.
- Cruz, B. A.; Pérez, A.; García, E.; Gallardo, F. y Soto, R. M. 2015. Análisis químico-morfológico comparativo de accesiones de *Jatropha curcas* L. del estado de Veracruz. Rev. Mex. Cienc. Agríc. 6(3):589-601.
- González, G. C. E. y Crespo, L. G. J. 2016. Response of *Moringa oleifera* Lam to fertilization strategies on lixiviated Ferralitic Red soil. Pastos y Forrajes. 39(3):173-177.
- Lok, S. y Suárez, Y. 2014. Efecto de la aplicación de fertilizantes en la producción de biomasa de *Moringa oleifera* y en algunos indicadores del suelo durante el establecimiento. Rev. Cub. Cienc. Agríc. 48(4):399-403.
- Medina, M.; García, D.; Clavero, T. e Iglesias, J. 2007. Estudio comparativo de *Moringa oleifera* y *Leucaena leucocephala* durante la germinación y la etapa inicial de crecimiento. Zootecnia Tropical. 25(2):83-93.
- Olivares, M. A.; Hernández, A.; Vences, C. y Jáquez, J. L. 2012. Lombricomposta y composta de estiércol de ganado vacuno lechero como fertilizantes y mejoradores de suelo. Universidad y Ciencia. 28(1):27-37.
- Oloyede, D.; Aviara, N. y Shittu, S. 2015. Measurement of engineering properties necessary to the design of drumstick (*Moringa oleifera* L.) Pod Sheller. J. Biosys. Eng. 40(3):201-211.
- Olson, M. y Fahey, J. 2011. *Moringa oleifera*: un árbol multiusos para las zonas tropicales secas. Revista Mexicana De Biodiversidad. 82(4):1071-1082.
- Pérez, A.; Sánchez, T.; Armengol, N. y Reyes, F. 2010. Características y potencialidades de *Moringa oleifera*, Lamark. Una alternativa para la alimentación animal. Pastos y Forrajes. 33(4):1-16.
- Pérez, R.; de la Cruz, J.; Vázquez, E. y Obregón, J. 2010. *Moringa oleifera*, una nueva alternativa forrajera para Sinaloa. C. Consultivo, Ed. Culiacán. Fundacion produce Sinaloa, AC. 1-29.
- Quintas, G.; Valdés, O. A. y Pérez, A. 2016. Efecto de la poda y fertilización orgánica en *Moringa oleifera* Lam. en la región centro de Veracruz, México. Rev. Cienc. Adm. 1(especial):101-121.
- Ramos, D. y Terry, E. 2014. Generalidades de los abonos orgánicos: importancia del Bocashi como alternativa nutricional para suelos y plantas. Cultivos Tropicales. 35(4):52-59.
- Restrepo, J. 2001. Elaboración de abonos orgánicos fermentados y biofertilizantes foliares Experiencias con agricultores en Mesoamérica y Brasil. Costa Rica. 155 p.
- Restrepo, J. y Hensel, J. 2006. El A, B, C de la agricultura orgánica y panes de piedra. México.
- Restrepo, J. y Hensel, J. 2009. Manual práctico de agricultura orgánica y panes de piedra. Primera ed. Colombia. 318 p.
- Rodríguez, M. M.; Soto, O. R.; Parets, S. E. y Alemán, P. R. 2005. Bocashi, una alternativa para la nutrición de la habichuela (*Vigna unguiculata* L. Walp sub-sp *sesquipedalis* L.), variedad Cantón 1 en huertos populares. Rev. Centro Agric. 32(1):71-76.

- Sarwar, M.; Ali, A.; Nouman, W.; Arshad, M. I. and Patra, J. K. 2017. Compost and synthetic fertilizer affect vegetative growth and antioxidant activities of moringa. *Inter. J. Agric. Biol.*
- Sosa, A. A.; Ledea, J. L.; Estrada, W. y Molinet, D. 2016. Efecto de la distancia de siembra en variables morfoagronómicas de moringa (*Moringa oleifera*). *Agron. Mesoam.* 28(1):207-211.
- Swati, B.; Kamal, S. and Dixit, B. 2013. Fertilizer management in drumstick under semidry condition of Bundelkhand region of Madhya Pradesh. *Nature Sci.* 11(2):67-68.
- Toral, O.; Cerezo, Y.; Reino, J. y Santana, H. 2013. Caracterización morfológica de ocho procedencias de *Moringa oleifera* (Lam.) en condiciones de vivero. *Pastos y Forrajes.* 36(4):409-416.
- Usman, D. C.; Usman, C.; Bonilla, R. y Sánchez, M. S. 2003. Efecto de la fertilización orgánica sobre la producción de follaje y rendimiento de semilla de cilantro *Coriandrum sativum* L. variedad Unapal Precoso. *Acta Agron.* 52(1-4):59-63.
- Valdés, O.; Palacios, O.; Ruíz, R. y Pérez, A. 2014. Potencial de la asociación moringa y ricinus en el subtrópico veracruzano. *Rev. Mex. Cienc. Agríc.* 9:1673-1686.
- Valdés, R. O. A.; Muñoz, G. C.; Pérez, V. A. y Martínez, P. L. E. 2014. Análisis y ajuste de curvas de crecimiento de *Moringa oleifera* Lam. en diferentes sustratos. *Rev. Biol. Agropec. Tuxpan.* 2(2):66-70.
- Velázquez, M.; Peón, I. E.; Zepeda, R. y Jiménez, M. A. 2016. Moringa (*Moringa oleifera* Lam.): usos potenciales en la agricultura, industria y medicina. *Rev. Chapingo Ser. Hortic.* 22(2):95-116.
- Zayed, M. S. 2012. Improvement of growth and nutritional quality of *Moringa oleifera* using different biofertilizers. *Annals Agric. Sci.* 57(1):53-62.