

Frequency of hypertension, age and gender in cerebrovascular disease in Paraguay

Luis Díaz-Escobar, Alan Flores*, Laia Seró-Ballesteros, Christian Otto, Ricardo Mernes, Fátima Pedrozo, Fabiola Riquelme, and Romina Gonzalez

Unidad de Ictus, Hospital de Clínicas, Servicio de Urgencias, Facultad de Ciencias Médicas, Universidad Nacional de Asunción, San Lorenzo, Paraguay

Abstract

Background: Hypertension, age, and gender are relevant factors associated to cerebrovascular disease. In Paraguay, cerebrovascular disease is the fourth cause of mortality; however, information about demographics and baseline characteristics is not widely knowledge. Our aim was determinate the prevalence of hypertension, age, and gender in the setting of acute-phase stroke in our population. **Methods:** This is a descriptive, single-center study. Data were collected from a prospective registry of stroke patients admitted in the stroke unit of our center, from April 2015 to December 2016. **Results:** From 996 stroke patients, 252 (25.3%) presented intracerebral hemorrhage (ICH). Three hundred ninety (39.1%) were female. The hypertension rate was 74% and 78% in ischemic stroke and ICH, respectively, being hypertensive hemorrhagic etiology 79.5% of the ICH and lacunar infarct 11.6% of ischemic strokes. In ICH patients, mean age differences between genders were remarkable (mean years 55.41 [SD \pm 14.8] vs. 62.48 [SD \pm 15.2] $p \leq 0.001$). In the multivariant analysis, lower age than 60 years old (OR: 4.893; CI 95%: 1.772-13.509 $p = 0.029$) and higher systolic blood pressure at admission (OR: 1.098; CI 95% 1.044-1.155, $p = 0.009$) were independent factors associated to ICH. **Conclusion:** In our population, ICH rates are similar to regional findings, occurring at an early age than other series, being remarkable in males. Hypertension rates in ischemic stroke and ICH are higher than other series, and the variability of presumed hypertensive microangiopathy phenotype could be in relation to age (hypertensive hemorrhage vs. lacunar). These findings would be related to ethnics, social, environment, and geographies factors.

Keywords: Stroke epidemiology. Hypertensive hemorrhage. Female stroke. Vascular risk factors. Intracerebral hemorrhage.

Frecuencia de la hipertensión arterial, la edad y el género en la enfermedad cerebrovascular en Paraguay

Resumen

Background: La hipertensión, la edad y el género son factores relevantes asociados a enfermedad cerebrovascular (ECV). En Paraguay, la ECV es la cuarta causa de mortalidad, sin embargo, la información sobre sus características no es de amplio conocimiento. Nuestro objetivo fue determinar la frecuencia de la hipertensión, la edad y el género en la ECV en fase aguda en nuestra población. **Métodos:** Se trata de un estudio observacional descriptivo monocéntrico. Se recogieron datos de

Correspondence:

*Alan Flores

E-mail: alanflo2507@hotmail.com

2604-6180 / © 2019 Academia Mexicana de Neurología A.C. Published by Permayer México. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 24-04-2019

Date of acceptance: 29-05-2019

DOI: 10.24875/RMN.19000078

Available online: 02-05-2022

Rev Mex Neuroci. 2022;23(3):80-85

www.revexneurociencia.com

un registro prospectivo de pacientes con ictus ingresados en la Unidad de Ictus de nuestro centro, desde abril de 2015 a marzo de 2019. **Resultados:** De 996 pacientes con ictus, 252 (25,3%) presentaron hemorragia intracerebral (HIC). Trescientos noventa (39.1%) fueron mujeres. La tasa de hipertensión arterial fue elevada tanto en isquémicos (74%) como en hemorrágicos (78%), siendo la hemorragia hipertensiva el 79.5% de las HIC y los lacunares, el 11.6% de ictus isquémicos. En pacientes con HIC, la diferencia de edad entre géneros fue significativa (media/años 55.41 (SD ± 14.8) vs. 62.48 (SD ± 15.2) $p \leq 0.001$). En el análisis multivariante, fueron factores independientes asociados a HIC: la edad menor a 60 años (OR: 4.893; IC 95%: 1.772-13.509 $p = 0.029$) y una elevada presión arterial sistólica al ingreso (OR: 1.098; IC 95% 1.044-1.155, $p = 0.009$). **Conclusión:** En nuestra población, la frecuencia de HIC es similar a lo descrito en la región, observándose una edad más temprana que en otras series y siendo remarcable en el género masculino. La tasa de hipertensión en HIC e ictus isquémico es mayor que en otras series y la variación de distintos fenotipos por presunta microangiopatía hipertensiva estaría relacionado en parte a la edad (hemorragia hipertensiva vs. lacunares). Estos resultados podrían explicarse por razones étnicas, socioambientales y geográficas.

Palabras clave: Ictus epidemiología. Hemorragia cerebral hipertensiva. Ictus en la mujer. Factores de riesgo. Hemorragia intracerebral.

Introduction

Stroke is one of the diseases with the greatest impact on public health, being the third cause of mortality and the leading cause of disability in the Western world since most patients suffer sequels, which is 30% of cases disable them to perform daily activities^{1,2}.

The proportion and frequency between the ischemic types and intracerebral hemorrhage (ICH) are variables according to a determined region; while in Spain, the rates of ICH are around to 12%³, in regions of South America such as Ecuador and Chile the ICH reach rates between 37 and 46%⁴. In Paraguay, cerebrovascular disease is the fourth cause of death⁵. However, another data about the baseline characteristics in our population are not widely known.

Among the cardiovascular risk factors related to stroke, hypertension is the most frequent and important risk factor. The prevalence is around 70% in both ischemic and hemorrhagic patients, and its optimal control is possibly one of the most important interventions in secondary prevention for both types of stroke^{3,6}. In Paraguay, the prevalence of hypertension is high, with rates of 53% in men and 40% in women between 35 and 45 years, reaching up to 81% in the general population between 65 and 74 years⁷.

Another known and controversial stroke risk factor is gender, and their disparities in baseline and clinical outcomes. Several descriptive studies have shown an association of older age, frequency of hypertension, obesity, and atrial fibrillation in women than in men^{8,9}.

Objective

The objective of the study was to determine the rates of hypertension, age, gender, and other known risk

factors in patients with stroke in the acute phase in our environment.

Methodology

This is a descriptive, observational study in a single center. Data were collected from a prospective of stroke patients admitted in the Stroke Unit of our center from April 2015 to March 2019.

The known modifiable and non-modifiable risk factors have been recorded. In patients with hemorrhagic stroke, the etiology of hypertensive ICH has been determined according to the history of hypertension, the topography of the parenchymal lesion, and the vascular study¹⁰. The diagnosis of ischemic stroke was made according to the criteria of the Oxfordshire Community Stroke Project and the Trial of ORG 10172 in Acute Stroke Treatment¹¹.

The descriptive statistical analysis was performed for all the parameters, which included measurements of the central tendency and dispersion for the quantitative variables, as well as absolute and relative frequencies for the qualitative variables, with their 95% CI in both cases. If the data did not meet the assumptions of normality for the analysis, non-parametric statistical methods were used (Student's t-test or Mann-Whitney test, respectively). The Chi-square test was applied for the comparison between the proportions. To determine the weight of each of the risk factors between ischemic and hemorrhagic stroke, multiple logistic regression models were established, with a 95% CI. The statistical tests were performed with a significance level of 5%. The statistical program SPSS v. 23.0 was used to perform the analysis.

Table 1. Baseline characteristics of all patients with stroke and subgroups; ischemic and hemorrhagic with their significance in the univariate analysis

	All (n = 996)	Ischemic (n = 744)	ICH (n = 252)	p-value
Age, year	62.8DS ± 15.3	64.3DS ± 15.2	58.1 ± 14.8	< 0.001
Glicemia at admission mgr/dl. (n:169)	165.4DS ± 120.3	168.6DS ± 97.7	155.9DS ± 120.3	0.513
Female gender n (%)	390 (39.1)	295 (39.6)	93 (36.9)	0.585
Hypertension n (%)	749 (75.2)	553 (74.3)	195 (77.3)	0.102
Diabetes mellitus n (%)	237 (23.6)	206 (23.5)	30 (3.4)	< 0.001
Dyslipidemia n (%)	56 (5.6)	47 (5.4)	9 (1)	0.144
Atrial fibrillation	128 (12.7)	117 (13.6)	11 (1.3)	< 0.001
Prior antiplatelet treatment	996 (99.1)	81 (8.1)	10 (1.0)	0.003
Prior stroke	150 (14.9)	129 (15.4)	21 (2.5)	0.001
Baseline NIHSS median (interquartile range)	8 (4)	8 (4)	9 (4)	0.796
Systolic blood pressure at admission	169.8 ± 38.3	161.4 ± 32.5	189.9 ± 43.2	< 0.001

Results

From 996 patients with stroke, 744 (74.7%) were ischemic, and 252 (25.3%) were ICH, and 390 (39.1%) were women. The mean age was 62.8 years (SD ± 15.3). Overall, 75.4% had known antecedent of hypertension, 27% DM, and 6.4% dyslipidemia. The median baseline NIHSS was 8 (IQR: 4-12). The baseline characteristics in general and of the ischemic and hemorrhagic groups are shown in [table 1](#).

In ischemic stroke patients, the mean age was 64.3 (SD ± 15.2) years, the median of NIHSS was 8 (IQR 4-12), and 39.8% were women. Patients with ischemic stroke were significantly older (mean years 64.3 SD ± 15.2 vs. 58.1 SD ± 14.8), had higher glycemia at admission (mean mgr./dl 168.6 SD ± 97.7 vs. 155.9 SD ± 120.3 p = 0.513), frequency of diabetes mellitus (23.5% vs. 3.4% p = 0.001), atrial fibrillation (13.6% vs. 1.3% p = 0.001), prior history of stroke (15.4% vs. 2.5% p = 0.001), prior antiplatelet treatment (8.1% vs. 1.0% p = 0.001), and lower systolic blood pressure on arrival than patients with ICH (mean mmHg, 169.8 SD ± 38.3 vs. 189.9 SD ± 43.2).

The most frequent etiology was (422 cases work-up performed) cardioembolic 41.2%, indeterminate in 24.1%, lacunar infarcts in 11.6%, and atherothrombotic in 21.5%. Hypertension was present in 91% of atherothrombotic strokes, 89.6% of lacunary strokes, 80.8% of cardioembolic, and 77.6% of indeterminate strokes ([Fig. 1](#)).

ICH patients had a mean age of 58.1 (SD ± 14.8) years, the median of NIHSS was 9 (IQR 5-13), and 36.9% were women. From 142 cases with etiological workup, 79.5% were due to hypertensive hemorrhage. The factors significantly related to hypertensive hemorrhage (113 cases evaluated) were hypertension (p ≤ 0.001) and the lower age stratified by groups (p = 0.006) ([Fig. 2](#)).

In the multivariate analysis adjusted for sex, hypertension, diabetes mellitus, previous stroke, and atrial fibrillation, age < 60 years old (OR: 4.893, 95% CI: 1.772-13.509, p = 0.029) and a higher systolic blood pressure at admission (OR: 1.098, 95% CI 1.044-1.155, p = 0.009) were independent factors associated with ICH.

In relation to gender, hypertension was present in 79.3% in men and 83.5% in women. In men, smoking habit was significantly more frequent (23.9% vs. 3.4% p < 0.001) and they presented more ischemic heart disease (7.9% vs. 2.3% p = 0.001) than women. Men were younger (61.75 SD ± 13.6 vs. 64.16 SD ± 17.4 p = 0.003), being in the ICH group, even more remarkable the difference (mean years 55.41 [SD ± 14.8 vs. 62.48 [SD ± 15.2 p ≤ 0.001) ([Fig. 3](#)).

Discussion

This is a descriptive study, with data from a prospective cohort over a period of 41 months. In general, there is a higher proportion of men than women with stroke,




Figure 1. Graphic bar showing the number of cases of ischemic stroke etiology according to prior hypertension status.




Figure 2. Graphic bar showing the number of cases of intracerebral hemorrhage according to stratified age group.

and the mean age is significantly lower than in developed countries^{1,3}, but similar to previous studies in

Buenos Aires and Brasilia^{12,13}. Hypertension is the most frequent risk factor, slightly higher than other series^{3,12-14}. Another risk factors rate such as DM, previous stroke, and atrial fibrillation were similar to prior studies; however, the frequency of dyslipidemia, known ischemic cardiopathy, and smoking was strikingly lower^{3,12-14}. Possible explanations (in addition to the inherent biases and limitations from the nature of this study) are that our center, although it is located in an urban center, is a reference center that receives patients from the rural area also, and this could affect our results. Another reason would be the lack of knowledge of the risk factors of our population (less than half of men know about their blood cholesterol levels)⁷.

The rate of HIC is significantly higher than that described in developed countries^{3,12} and is consistent with results from our region in studies conducted in Chile, Ecuador, and Argentina¹⁴⁻¹⁶. These results are explained by the high prevalence rate of hypertension in our population, especially manifests at early age in adult⁷.




Figure 3. Bars showing the mean age according to gender in overall **A:** ischemic; **B:** intracerebral; and **C:** stroke patients.

In relation to ischemic stroke, the etiology by TOAST (in 422 cases) showed a high rate of cardioembolic (41.2%), however a lower frequency of atheromatous disease (21.5%) and lacunar (11.6%). These results are more similar to those found in Santiago¹⁴ than in those shown in Brasilia, Guayaquil, and Buenos Aires^{13,15,16}. It should be noted that in South America the frequency in relation to the etiology of stroke is very heterogeneous and diverse, mainly due to ethnic, geographical, and sociocultural differences⁴. Finally, the relatively high rate of indeterminate etiology would be related in many cases to an incomplete study and is a clear limitation in the present study.

Interestingly, while hypertension rates are high (79.3%) in ischemic patients, the percentage of lacunar strokes for presumed hypertensive microangiopathy as a cause is lower than expected (11.6%). In contrast, hypertensive hemorrhage represents the cause of up to 79.5% of ICH. Previous studies in relation to this possible common substrate have shown that the cerebral vascular phenotype (ICH vs. lacunar infarcts) due to hypertensive microangiopathy would be related in part to age and cholesterol levels^{17,18}. Therefore, a possible explanation to our findings could be related to age (remarkably younger patients) and the lower frequency of dyslipidemia (5.6%) known in our population. Another possible explanation would be in relation to the less severe clinical deficit associated with lacunar stroke¹⁹, which may affect the arrival or referral of these patients. These results are consistent in the region with previous studies in Santiago de Chile and Bogotá^{13,20}. Unfortunately, the body mass index, which has been shown to be a prognostic factor for the development of ICH or lacunar infarctions in hypertensive microangiopathy¹⁸, has not been evaluated in the present study. Future prospective and comparative studies should be taken into account to confirm these findings.

Regarding gender, women presented a stroke less frequently, with greater age (according to the

literature)^{8,21,22}, and significantly less known cardiovascular risk factors than men, which is also consistent with previous studies. Factors such as hypertension and atrial fibrillation, which in other series showed a significantly higher frequency in women, have not been seen in our study^{8,9}. The age difference in ICH is remarkable among genders, being the higher hypertension prevalence in early age⁷ and the hypothetical hormonal preventive effect in women^{20,21} probable reasons for this finding.

As a relevant limitation at the present study; it is a monocentric study, involving patients from urban and rural areas that may not represent a specific geographic area.

Conclusion

In this prospective study in our country, hypertension is a risk factor highly related to stroke with a higher frequency of hypertensive hemorrhages in younger patients than other series, especially in men. The frequency of different stroke subtypes would be related to environmental, sociocultural, and biological factors. Multicenter and comparative studies are warranted to confirm these findings. The intensification of public health programs focused on the control of risk factors, especially hypertension, is necessary to prevent cerebrovascular disease in our environment.

Funding

There was no particular funding source for this scientific report.

Conflicts of interest

The authors declare that in this study, there are no relevant conflicts of interest.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

References

- Bonita R. Epidemiology of stroke. *Lancet*. 1992;339:342-4.
- Truelsen T, Piechowski-Józwiak B, Bonita R, Mathers C, Bogousslavsky J, Boysen G, et al. Stroke incidence and prevalence in Europe: a review of available data. *Eur J Neurol*. 2006;13:581-98.
- Arias-Rivas S, Vivancos-Mora J, Castillo J, En Nombre de Los Investigadores Del Registro Epices. Epidemiology of the subtypes of stroke in hospitalised patients attended by neurologists: results of the EPICES registry (I). *Rev Neurol*. 2012;54:385-93.
- Saposnik G, Del Brutto OH, Iberoamerican Society of Cerebrovascular Diseases. Stroke in South America: a systematic review of incidence, prevalence, and stroke subtypes. *Stroke*. 2003;34:2103-7.
- Boletín de Vigilancia. Enfermedades No Transmisibles Y Factores De RIESGO 2015. Ministerio de Salud Pública y Bienestar Social. Available from: <http://www.mspbs.gov.py>.
- Kernan WN, Ovbiagele B, Black HR, Bravata DM, Chimowitz MI, Ezekowitz MD, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American heart association/American stroke association. *Stroke*. 2014;45:2160-236.
- Primera Encuesta Nacional de Factores de Riesgo de Enfermedades No Transmisibles. Primera Edición. Asunción. 2012 Ministerio de Salud Pública y Bienestar Social. Available from: <http://www.mspbs.gov.py>.
- Andersen KK, Andersen ZJ, Olsen TS. Age-and gender-specific prevalence of cardiovascular risk factors in 40,102 patients with first-ever ischemic stroke: a nationwide Danish study. *Stroke*. 2010;41:2768-74.
- Reeves MJ, Bushnell CD, Howard G, Gargano JW, Duncan PW, Lynch G, et al. Sex differences in stroke: epidemiology, clinical presentation, medical care, and outcomes. *Lancet Neurol*. 2008;7:915-26.
- Lang EW, Ren Ya Z, Preul C, Hugo HH, Hempelmann RG, Buhl R, et al. Stroke pattern interpretation: the variability of hypertensive versus amyloid angiopathy hemorrhage. *Cerebrovasc Dis*. 2001;12:121-30.
- Bamford J, Sandercock P, Dennis M, Burn J, Warlow C. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet*. 1991;337:1521-6.
- Saposnik G, Caplan LR, Gonzalez LA, Baird A, Dashe J, Luraschi A, et al. Differences in stroke subtypes among natives and Caucasians in Boston and Buenos Aires. *Stroke*. 2000;31:2385-9.
- Carod-Artal FJ, Casanova Lanchipa JO, Cruz Ramírez LM, Pérez NS, Siacara Aguayo FM, Moreno IG, et al. Stroke subtypes and comorbidity among ischemic stroke patients in Brasília and Cuenca: a Brazilian-spanish cross-cultural study. *J Stroke Cerebrovasc Dis*. 2014;23:140-7.
- Nogales-Gaete J, Núñez L, Arriagada C, Sáez D, Figueroa T, Fernández R, et al. Caracterización clínica de 450 pacientes con enfermedad cerebrovascular ingresados a un hospital público durante 1997. *Rev Med Chil*. 2000;128:1227-36.
- Del Brutto OH, Mosquera A, Sánchez X, Santos J, Noboa CA. Stroke subtypes among hispanics living in Guayaquil, Ecuador. Results from the Luis Vernaza hospital stroke registry. *Stroke*. 1993;24:1833-6.
- Saposnik G, Gonzalez L, Lepera S, Luraschi A, Sica RE, Caplan LR, et al. Southern Buenos Aires stroke project. *Acta Neurol Scand*. 2001;104:130-5.
- Labovitz DL, Boden-Albala B, Hauser WA, Sacco RL. Lacunar infarct or deep intracerebral hemorrhage: who gets which? The Northern Manhattan study. *Neurology*. 2007;68:606-8.
- Lioutas VA, Beiser A, Himali J, Aparicio H, Romero JR, DeCarli C, et al. Lacunar infarcts and intracerebral hemorrhage differences: a nested case-control analysis in the FHS (Framingham heart study). *Stroke*. 2017;48:486-9.
- Pantoni L. Cerebral small vessel disease: from pathogenesis and clinical characteristics to therapeutic challenges. *Lancet Neurol*. 2010;9:689-701.
- Peña I, Ruiz C, Morillo LE, Pedraza OL, Sánchez E, Santín LC, et al. Clasificación del TOAST en la práctica clínica de un hospital universitario. *Acta Neurol Colomb*. 2001;17:304-8.
- Alonso de Leciana M, Egidio JA, Fernández C, Martínez-Vila E, Santos S, Morales A, et al. Risk of ischemic stroke and lifetime estrogen exposure. *Neurology*. 2007;68:33-8.
- Murphy SJ, McCullough LD, Smith JM. Stroke in the female: role of biological sex and estrogen. *ILAR J*. 2004;45:147-59.